

Numerical Simulations of Astrophysical Jets from Keplerian Disks with Periodic Ejection.

B. F. Vitorino, V. Jatenco-Pereira and R. Opher

Instituto de Astronomia, Geofísica e Ciências Atmosféricas

Universidade de São Paulo

Rua do Matão 1226, 05508-090 São Paulo, SP, Brazil

braulo@astro.iag.usp.br, jatenco@astro.iag.usp.br, opher@astro.iag.usp.br

Received _____; accepted _____

ABSTRACT

We present 2.5-dimensional time-dependent simulations of nonrelativistic and nonradiative outflows from sinusoidally perturbed Keplerian accretion disks. A sinusoidal perturbation is introduced in the velocity of the gas ejected from the surface of the disk into a cold corona. In the simulations, the disk is a fixed boundary from which the gas is ejected with a pulsed velocity. The maximum value of this velocity is taken to be a thousandth of the local Keplerian disk velocity. It was found that for large periods, the structures in the jet tend to fragment into smaller substructures. For small values of the period, the structures tend to dissipate, while for medium values of the period, they tend to persist.

Subject headings: ISM: jets and outflows—galaxies: jets—accretion, accretion disks—MHD.

1. Introduction

In general, the production of astrophysical jets has been primarily explained by two possible mechanisms: magnetic forces associated with an accretion disk (Königl & Pudritz 1999) and the interaction between an accretion disk and a magnetized star, in the case of protostellar jets (Shu et al. 1999). The first mechanism is a disk-wind driven model, where the gas is centrifugally flung out from the surface of a Keplerian disk when field lines thread the disk at an angle of 60° or less with respect to the surface. When the gas reaches the local Alfvén velocity, the $(\mathbf{J}_p \times \mathbf{B}_\phi)$ term becomes dominant, providing the collimation for the outflow. The second mechanism is the X-wind model, where the wind emanates near the region where the disk corotates with the central magnetic star. It is assumed that some magnetospheric lines, connecting the central star and the disk, are partially open.

Excess angular momentum brought in by the disk is extracted by the magnetocentrifugally driven wind. The wind emanates from the inner edge of the disk and flows out along the open magnetic field lines. Since the first mechanism seeks to explain astrophysical jets independently of the detailed nature of the central object, from active galactic nuclei (AGNs) to young stellar objects (YSOs), it has the possibility of being a universal model.

Recent theories and models of astrophysical jets have mainly tried to provide a natural explanation for common features. These include the episodic outbursts and knots that are observed.

Numerical simulations of axisymmetric magnetized flows from accretion disks have previously been made using different models. Shibata & Uchida (1986) and Stone & Norman (1994) followed the internal dynamics of the disk. Bell & Lucek (1995) and Ustyugova et al. (1995) treat the disk as a boundary condition at the base of the wind, without following its internal dynamics, and assume it to be in Keplerian equilibrium. Following the latter model, Ouyed & Pudritz (1997a, b; 1999) made 2.5-dimensional, high-resolution, numerical magnetohydrodynamic (MHD) simulations of the onset and collimation of outflows from the surface of a Keplerian accretion disk around a central object. Using two different topologies for the magnetic field that threads the disk and the corona, they obtained steady or episodic jets, depending on the values of the constant (in time) ejection velocity from the disk to the corona. For the smallest values of the ejection velocity, their results showed episodic production of knots near the surface of the disk which propagate along the jet axis. They argued that these knots arose due to MHD shocks.

Krasnopolsky et al. (1999) made time-dependent, axisymmetric simulations of the magnetocentrifugal model jet formation. Their system consisted of a Keplerian disk and a central object. The main subtlety resides in the treatment of the disk-wind boundary, the disk surface. Some, but not all, of the flow variables can be fixed on this boundary. The boundary

conditions imposed on the disk surface were limited to those necessary and sufficient to take into account information propagating upstream from the fast and Alfvén critical surfaces, avoiding overdetermination of the flow and the production of impulsive accelerations. They obtained a cold and steady jet, launched smoothly from the Keplerian disk and collimated by the magnetocentrifugal mechanism. The collimation is observed both in the shape of the field lines and in the density profiles, which become cylindrical, and thus jetlike, at large distances along the rotation axis. The structure of the flow is insensitive to the density or flow speed at the injection surface, as long as the mass flux is kept constant. As explained by the authors, steady-state jets obtained in earlier studies, using an essentially incomplete treatment of the disk boundary conditions, are qualitatively unchanged when this deficiency is rectified. It remains to be shown, however, whether the same is true for nonsteady jets.

Frank et al. (2000) made a series of simulations intended to address the issue of MHD jet propagation. Whereas some studies have used ad hoc initial conditions, they injected flows into a computational grid derived from models of collimated jets driven by magnetocentrifugal launching. The initial configurations in the jet were taken directly from the solution of the force balance perpendicular (the Grad-Shafranov equation) and parallel (the Bernoulli equation) to the magnetic surfaces generated by a magnetized rotator. Their simulations followed the evolution of jets composed of helical fields, embedded in hypersonic plasmas whose density and velocity varied with radius. They studied the propagation of jets driven by: (1) a purely Keplerian rotator; and (2) a Keplerian rotator with a sub-Keplerian boundary layer. Both rotators are exterior to a solid body rotator. Studying the adiabatic case for the last model (their multicomponent model) they obtained a jet in which strong instabilities developed in the core of the jet. Once the core is exposed, periodic pinches can be seen in the beam. Their stability analysis indicates that the origin of these periodic pinches could be current driven instabilities. The pinching instabilities on the axis have a wavelength of approximately half of the jet radius. However, the beam also has a second characteristic

wavelength which runs along the surface of the core. This feature, which appears as an envelope encompassing the shorter wavelength modes, has a wavelength of approximately three jet radii. As explained by the authors, these results suggest that these instabilities could be at the origin of the knotty structure of a large number of jets as seen, for example, in HL Tau, HH1, HH30, and HH34.

Time-dependent MHD calculations of a modified version of the X-wind model, considering the star-disk field interaction, shows twisting of closed stellar field lines, current sheet formation, reconnection, and ejection of magnetic island with the plasma heated to $\leq 10^8$ K, consistent with X-ray flare observations (Hayashi et al. 1996; Goodson et al. 1997).

All the above previous studies assumed constant velocity injection from the disk. Here, we extend the previous investigations and study the effect of non-steady injection, in particular, periodic injection.

Adopting the disk as a boundary condition, we investigated how periodic perturbations at the disk surface can alter the formation and the propagation of the magneto-centrifugal jet. In the FU-Orionis phase of a protostar, for example, we know that there is a periodic ejection of material observed in the X-ray range. We do not include the physics of the disk that could generate a pulsed gas ejection. We assume that there is a periodic mass loading at the base of the corona of the disks, which introduces a periodic ejection of gas from the disk surface. Whereas Ouyed & Pudritz (1997a, b; 1999) used a constant ejection velocity from the disk, we study the temporal dependence of this velocity. We made 2.5-dimensional numerical simulations using the ZEUS-3D code (Stone & Norman 1992a,b) in the axisymmetry option. A sinusoidal perturbation was introduced in the ejection velocity as a temporal function. Various values for the periods in the range from 10 to 80 rotation periods of the innermost radius of the accretion disk (r_i), were used in the simulations. The maximum value for the perturbation was chosen. This value is roughly equal to that which

would produce a steady state jet if the gas were continuously ejected from the disk (Ouyed & Pudritz 1997a, hereafter OP97a). The luminosity of the protostellar T Tauri stars (which have accretion disks) fluctuates rapidly with large amplitude. Similarly, quasars, which are also assumed to have accretion disks, also have luminosities which fluctuate rapidly with large amplitude. Therefore, we introduced in our simulations velocity perturbations that are a large percentage of the injection velocity. A region near the central object, $80r_i$ in the z -direction and $20r_i$ in the r -direction, was simulated.

Using different values for the period of the sinusoidal perturbation, we obtained the formation of regularly spaced structures along the jet axis. For small values of the period, the structures tend to dissipate along the jet axis; for medium values, they tend to persist and for large values, tend to fragment into smaller substructures.

In §2 the basic equations that are solved are presented. The model is described in §3 and the numerical results are presented and discussed in §4. Finally, in §5 we present a summary and the final conclusions.

2. Basic equations

Adopting cylindrical coordinates (r, ϕ, z) , we placed the central object at the origin and took the z -axis to be perpendicular to the disk. The surface of the disk thus lies in the $z = 0$ plane of our coordinate system.

The equations that describe the conservation of mass, momentum, energy and magnetic flux are:

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0, \tag{1}$$

$$\rho \left[\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} \right] + \nabla (p + p_A) + \rho \nabla \Phi - \mathbf{j} \times \mathbf{B} = 0, \quad (2)$$

$$\rho \left[\frac{\partial e}{\partial t} + (\mathbf{v} \cdot \nabla) e \right] + p (\nabla \cdot \mathbf{v}) = 0, \quad (3)$$

$$\frac{\partial \mathbf{B}}{\partial t} - \nabla \times (\mathbf{v} \times \mathbf{B}) = 0, \quad (4)$$

$$\nabla \cdot \mathbf{B} = 0, \quad (5)$$

with ρ the gas density, $\mathbf{v}$ the velocity, $\mathbf{B}$ the magnetic field, p the gas pressure, e the internal energy, $\mathbf{j} = \nabla \times \mathbf{B}/4\pi$ the electric current density, and $\Phi = -GM/(r^2 + z^2)^{1/2}$ the gravitational potential.

A polytropic gas $p = K\rho^\gamma$ with the polytropic index $\gamma = 5/3$ is assumed. Thus, ignoring radiative transfer effects, we do not solve the energy equation (3). We also assume that p_A is an Alfvénic turbulent pressure. This provides, together with the thermal pressure, the main support against gravity for the cold corona. Here $p_A = p/\beta_T$, where β_T is the ratio of the gas pressure to the Alfvénic turbulent pressure, assumed to be constant (OP97a).

All parameters are given in units of their values at the innermost radius of the accretion disk r_i . Thus, the normalized equation of motion (eq. [2]) is

$$\left[\frac{\partial \mathbf{v}'}{\partial t} + (\mathbf{v}' \cdot \nabla') \mathbf{v}' \right] = -\frac{1}{\delta_i} \frac{\nabla' (p' + p'_A)}{\rho'} - \nabla' \Phi' + \frac{2}{\delta_i \beta_i} \frac{\mathbf{J}' \times \mathbf{B}'}{\rho'}, \quad (6)$$

where the dimensionless variables are: $r' = r/r_i$, $z' = z/r_i$, $v' = v/v_{K,i}$, $\rho' = \rho/\rho_i$, $p' = p/p_i$, $B' = B/B_i$, $\Phi' = -1/(r'^2 + z'^2)^{1/2}$ and $\nabla' = r_i \nabla$. The parameter $\delta_i = \rho_i v_{K,i}^2 / p_i$ is the ratio of the Keplerian to the thermal energy density and $\beta_i = 8\pi p_i / B_i^2$ is the ratio of the gas to the magnetic pressure in the corona at r_i . At r_i , the Keplerian speed is $v_{K,i} = (GM/r_i)^{1/2}$

and B_i is the poloidal field at r_i . For convenience, we drop the primes and refer only to the normalized variables. Time is given in units of $t_i = r_i/v_{K,i}$, so that the dimensionless time is

$$\tau = \frac{t}{t_i}. \quad (7)$$

3. The Model

In this model, the disk is a fixed boundary condition in pressure equilibrium with the corona. At the center there is a point mass that represents a star or a black hole. An initial potential (zero current) poloidal field threads the disk and the corona.

The computational domain defines the spatial region where the MHD equations are time evolved. ZEUS-3D, a grid code, divides the computational domain into cells, called “active” zones. Finite differencing the evolution near the boundaries of the grid requires that values for the dependent variables be specified beyond the computational domain. Therefore, at each boundary, “ghost” zones are added. Values for the dependent variables in the ghost zones are specified, using boundary conditions appropriate to the geometry and the physics of the problem being solved. Thus, the evolution equations are not solved for the ghost zones.

We use inflow boundary conditions at the disk surface and outflow boundary conditions on the remaining boundaries. Reflecting boundaries are used along the axis of symmetry, which coincides with the disk axis. The normal component (radial component) of the velocity and the magnetic field are set to zero on the boundary and reflected into the “ghost” zones. In addition, the 3-component (toroidal component) of the velocity and magnetic field is also inverted along the axis of symmetry into the “ghost” zones. The initial setup and boundaries are shown in Figure 1 (Fig. 2 of OP97a).

EDITOR: PLACE FIGURE 1 HERE.

In the corona, the initial toroidal field is zero. Magnetostatic equilibrium is numerically maintained to assure that an outflow can only arise due to the magnetocentrifugal mechanism. Resolving equation (6) for magnetostatic equilibrium, we obtain the expression for the coronal density distribution,

$$\rho = \left(1/\sqrt{r^2 + z^2}\right)^{3/2}, \quad (8)$$

which is used in our numerical simulations.

Since the surface of the accretion disk is a fixed boundary condition, there is no back reaction by the jet. We are, thus, time evaluating the coronal region ($z \geq 0$) only where the jet arises. However, we fix the density at the base of the corona (first column of cells), assuming that the disk is a suitable dynamic source of matter. The density distribution of the disk surface has the form

$$\rho_d = \eta_i r^{-3/2}, \quad (9)$$

where $\eta_i = \rho_d/\rho_0$ is the ratio between the disk density (ρ_d) and the density at the base of the corona ($\rho_0 \equiv \rho(z = 0)$).

The corona and the disk are threaded by the poloidal magnetic field in a potential configuration ($\mathbf{J} = 0$), described by the stream function,

$$\Phi_m = \sqrt{r^2 + (z_d + z)^2} - (z_d + z), \quad (10)$$

where z_d is the dimensionless disk thickness. The field components are given as

$$B_z = \frac{1}{r} \frac{\partial \Phi_m}{\partial r} \quad \text{and} \quad B_r = -\frac{1}{r} \frac{\partial \Phi_m}{\partial z}. \quad (11)$$

Due to the torsion of the poloidal field at the disk surface, a toroidal magnetic field is produced at the base of the corona, which could propagate into the disk. We thus expect that the disk has a nonzero toroidal field,

$$B_\phi = \frac{\mu_i}{r}, \quad (12)$$

where the input parameter $\mu_i = B_{\phi i}/B_i$ is the ratio of the toroidal to poloidal magnetic field at r_i . There are three important initial timescales defined at the disk's surface: the Kepler time τ_K , the Alfvén time τ_A and the magnetic braking time τ_B . Thus, we have (e.g., eq. [4.64] of OP97a)

$$(\tau_K, \tau_A, \tau_B) = \left(1, \sqrt{\delta_i \beta_i / 2}, \eta_i \sqrt{\delta_i \beta_i / 2}\right). \quad (13)$$

To estimate the turbulent pressure needed to support the corona for a given gas pressure, we use the equation for the ratio of the turbulent to magnetic pressure β_T

$$\beta_T = \gamma / ((\gamma - 1) \delta_i - \gamma), \quad (14)$$

where, as in equation (6), δ_i is the ratio of the Kepler to thermal energy density. The gas is injected from the disk into the corona at a velocity $\mathbf{v}_p = f_v v_K \mathbf{B}_p / B_p$, where $v_K = (1/r)^{1/2}$ is the local Keplerian velocity at the disk surface (for $r \geq 1$)¹, $\mathbf{B}_p$ is the local poloidal magnetic field that threads the disk, and f_v is the temporal function that describes the temporal dependence of the injection velocity. In our simulation we introduced a pulsed injection velocity with a sinusoidal form,

$$f_v = v_{inj0} [1 + \sin(2\pi\tau/T)], \quad (15)$$

where T is the period of the perturbation, v_{inj0} is an input parameter and τ is given by equation (7).

We, thus, have six free parameters in our simulation: δ_i , β_i , η_i , μ_i , v_{inj0} and T .

¹For the radial region where the disk does not exist ($r < 1$ (i.e., $r < r_i$)) we set $\mathbf{v}_p$ to zero.

4. Numerical results and discussion

We used the following parameters to investigate the time evolution of a jet from a Keplerian accretion disk with pulsed gas injection (similarly to OP97a):

$$(\delta_i, \beta_i, \eta_i, \mu_i, v_{inj0}) = (100.0, 1.0, 100.0, 1.0, 10^{-3}), \quad (16)$$

implying the initial timescales

$$(\tau_K, \tau_A, \tau_B) = (1.0, 7.07, 707.1). \quad (17)$$

Five simulations were made. In simulation A, there was no perturbation; in simulations B to E, we introduced perturbations with various values for the period T . The simulations were made in the domain $(z, r) = (80, 20)$ with a resolution of $(120, 45)$ for 400 time units ($< \tau_B$). Along the z-axis, 30 zones span a distance of $10r_i$ outwards from the disk surface. An additional 90 ratioed zones span an additional distance of $70r_i$, with each subsequent zone increasing in size by a factor of 1.017. Twenty uniform zones span an initial five disk radius in the radial direction. Outside of these uniform grid zones, 25 zones are ratioed, with each subsequent zone increasing in size by a factor 1.065.

Figure 2 shows the initial conditions in the atmosphere. The density distribution (eq. [8]) is shown at the top and the initial magnetic (potential) configuration, at the bottom of the figure. Initially, the velocity and the toroidal magnetic field are zero.

EDITOR: PLACE FIGURE 2 HERE.

The results of the simulations are shown in a series of figures that are described below. Figures 3 to 7 show the distributions of density, poloidal velocity (vectors), poloidal and toroidal magnetic fields and toroidal velocity at 400 time units for the simulations A to E, respectively. The contour lines are isocontours in the $r - z$ plane.

In simulation A (Fig. 3), no perturbation was introduced in the gas injection. Results in good agreement with those of OP97a were obtained. From the density distribution, shown at the top of this figure, we observe the formation of a highly collimated steady state jet along the z-axis. The collimation is due to the strong radial pinch force of a dominant toroidal field (fourth panel from top to bottom) that propagates into the corona. We obtain a maximum value for the poloidal velocity (second panel from top to bottom) on the order of the Keplerian velocity at r_i .

EDITOR: PLACE FIGURE 3 HERE.

A sinusoidal perturbation in the gas injection velocity (eq. [15]) at the disk surface was introduced in the other four simulations. As we are interested in studying the behaviour of the jet subject to this perturbation at its base, we chose a progressive range for the period T . In simulations B to E, we use $T = 10, 20, 40, 80$, respectively.

Along the jet axis in simulation B for $T = 10$ (Fig. 4), we observe the formation of regularly spaced structures up to about $z = 50r_i$. The distance of separation between two consecutive structures is about $5r_i$ (i.e., $\sim (T/2)r_i$). The axial extent of the regularly spaced structures advance with the flow speed until it reaches the limiting distance of about $50r_i$. This pattern persists even for times greater than 400 time units.

EDITOR: PLACE FIGURE 4 HERE.

Our results were obtained using ratioed zones that allow for a smaller number of calculated zones and, therefore, a poorer resolution as compared with OP97a. To verify the sensitivity of our results to this poorer resolution, we made simulation C using only uniform zones at a resolution close to that used in the uniform zone region in the poorer resolution

case. This simulation was run in the domain $(z, r) = (80, 20)$ with the resolution of $(250, 80)$ uniform zones. For the others cases (D – E), we used the poorer resolution.

Regularly spaced structures along the whole jet (Fig. 5) are seen in simulation C, where $T = 20$. These structures persist during the entire simulation, even for times greater than 400 units. The distance of separation between two consecutive structures is about $11r_i$, or roughly $(T/2)r_i$. We obtained these same results when using ratioed zones in the poorer resolution case.

EDITOR: PLACE FIGURE 5 HERE.

We observe the formation of regularly spaced structures along the entire jet as well as their fragmentation in simulation D for $T = 40$. Few minor structures are observed between the major structures (Fig. 6: top). Fragmentation of the structures can be seen in the isocontours of the toroidal velocity in the bottom panel of Figure 6. These structures persist during the entire simulation, even for times greater than 400 units. The distance of separation between two major structures is about $22r_i$, i.e., roughly $(T/2)r_i$.

EDITOR: PLACE FIGURE 6 HERE.

In simulation E for $T = 80$, we observe the formation of regularly spaced structures along the entire jet (Fig. 7: top), with more pronounced fragmentation clearly seen in the plot of the toroidal velocity (Fig. 7: bottom). These structures persist during the entire simulation, even for times greater than 400 units. The distance of separation between two major structures is about $40r_i$, i.e., $\sim (T/2)r_i$.

EDITOR: PLACE FIGURE 7 HERE.

The axial distributions, for all the simulations, of density, axial velocity, toroidal velocity and toroidal magnetic field along a cut at $r = 5r_i$, parallel to the disk’s axis at 400 time units, are shown in Figure (Fig. 8). These quantities are normalized to their maximum values along this cut. We note from the axial velocity distribution (dotted line), that the flow is rapidly accelerated along the field line from the disk to roughly $z = 15r_i$. After this point there is a “Hubble flow” constant acceleration, with v_z proportional to z for simulation A. However, for simulations B to E this constant acceleration character is modified, reflecting the pulsed nature of the jet. From the toroidal magnetic field distribution (dashed-dotted line) we observe the effect of torsion of the poloidal field lines due to the rotation of the disk. The regular structures, in simulations B – E, have their minimum mass concentration (solid line) at the maximum values of toroidal magnetic field (dashed-dotted line), reflecting jet constriction by this field. The shift between the toroidal magnetic field and the density in the pinched structures is a general result of long standing (e.g., Chan & Henriksen (1980)) and it has been associated with the periodic oscillations caused by the pinch instability due to the toroidal magnetic field.

EDITOR: PLACE FIGURE 8 HERE.

For simulations A to E at $\tau = 400$, the axial and radial mass fluxes are shown in Figure 9. The mass flux is defined as

$$\dot{m} = \int \rho \mathbf{v} \cdot d\mathbf{A}, \quad (18)$$

where A is the area. The axial mass flux² refers to the quantity of mass per unit time which passes through a circular area $A = \pi r^2$ and is calculated for each z , using $r = 20r_i$;

²The computation of the axial mass flux involves computing the product of v_z and the area of each annulus for each radial zone $[\pi(r_{j+1}^2 - r_j^2)]$, where j is an active zone and r_j refers

the radial mass flux³ refers to the quantity of mass per unit time which passes through the area $A = 2\pi rz$ and is shown for each z , using $r = 20r_i$. These fluxes, given in units of $\dot{m}_i = 2\pi r_i^2 \rho_i v_{Ki}$, are important for the detection of the decollimation of the jet. Their values agree with the results of OP97a. At the time $\tau = 400$, the axial fluxes at $z = 80r_i$ are comparable to those at $z \sim 0$; the small difference owing to decollimation of the jet. As the axial mass flux decreases, the radial flux can be seen to increase by roughly the same amount. We can see the pulsed nature of the jet (simulations B to E) from the axial mass fluxes (solid lines) which oscillate around their average values. The radial mass fluxes in simulations B to E are similar to those of simulation A (without perturbation). In simulation B, the axial mass flux curve is not very modified by the density structures, as compared with the case for which there is no perturbation (simulation A); in simulations C to D, the axial mass flux curves are strongly modified by the density structures. Their average values, however, do not change.

EDITOR: PLACE FIGURE 9 HERE.

Figure 10 shows the sonic, Alfvénic and magnetosonic Mach numbers for all the simulations at $\tau = 400$, in a cut at $r = 5r_i$ along the z -axis. In the computation of the sonic Mach numbers, we use only the thermal pressure p in the sound speed. To compute the magnetosonic Mach number, we use the effective sound speed which includes the thermal

to the position of the face or edge of the zone. At each z , the axial mass flux is computed from the sum of the axial mass flux contributions of all 45 radial zones.

³The radial mass flux at each z involves computing the product of v_r and the area of the annulus for a given radial zone $[2\pi r_j(z_{k+1} - z_k)]$, where j and k define each active zone, and r_j and z_k refers to the face/edge position of the zone, along the r and z direction, respectively. In particular, we computed the radial flux at each z using $r = 20r_i$.

and the Alfvénic turbulent pressure, p_A ⁴. We note in Figure 10 that the jet is already supersonic and super-Alfvénic near the base of the corona and super-magnetosonic at $z \sim 4r_i$. Only where density structures exist are the Mach numbers (Fig. 10) modified. Their median values, however, remain roughly the same. The maximum values of Alfvénic Mach numbers (dashed-dotted line) are due to the decreased axial magnetic fields. Negative values correspond to a reversal of the axial magnetic field.

EDITOR: PLACE FIGURE 10 HERE.

5. Summary and Conclusions

We made numerical 2.5-dimensional simulations of astrophysical jets produced from Keplerian disks subject to a sinusoidal perturbation of the gas injection velocity from the disk into the corona. Ratioed zones were used in order to obtain better resolution near the region of the generation of the jet with a poorer resolution outside.

Five simulations were made with different values of the period of the sinusoidal perturbation. No perturbation was introduced in simulation A. In simulations B to E, we used periods $T = 10, 20, 40$ and 80 , respectively, in units of $t_i = r_i/v_{Ki}$, where r_i is the innermost radius of the disk and v_{Ki} , its Keplerian velocity.

For simulation A, where there was no perturbation, we obtained a steady state highly collimated jet with a maximum axial velocity of the order of v_{Ki} after a time $\tau = 400$. In a cut at $r = 5r_i$ parallel to the disk’s axis, the jet is supersonic and already super-Alfvénic

⁴The magnetosonic speed is defined as $(c_s^2 + v_{Az}^2)^{1/2}$ where $c_s = (\gamma(p + p_A)/\rho)^{1/2}$ is the effective sound speed and $v_{Az} = B/(4\pi\rho)^{1/2}$ is the axial Alfvén velocity.

near the base of the corona and super-magnetosonic at $z \sim 4r_i$. These results agree with those of OP97a, who used better resolution.

Introducing a sinusoidal perturbation in the gas injection velocity, the results common, to simulations B to E, are:

1. The formation of regularly spaced structures with a separation distance of roughly $(T/2)r_i$ along the jet axis.
2. The toroidal magnetic field is out of phase ($\sim 90^\circ$) with the density curve along the jet axis.
3. The axial mass fluxes are modified by the periodic density structures, although the average remains roughly the same, as compared with the same curve for the case of no perturbation (simulation A). The axial fluxes at $z = 80r_i$ are comparable to the axial fluxes at $z \sim 0$.
4. The radial mass fluxes are similar to those of simulation A (where there was no perturbation). However, differences arise when we examine a cut at $r = 5r_i$. There is oscillatory behaviour in some of the simulations at this cut.
5. In a cut at $r = 5r_i$ the Mach number distributions are modified only where density structures exist. The median values, however, remain roughly the same, and are similar to simulation A.

The results that are different in some of these simulations are:

- a. In simulation B ($T = 10$) the structures persist only up to an axial distance of $z \simeq 50r_i$.
- b. In simulations D ($T = 40$) and E ($T = 80$), fragmentation of the structures occur.

These simulations, introducing periodic perturbations in the accretion disk, produced

collimated jets with a wavelength of $(T/2)r_i$, where T is the temporal period of the perturbation. The creation of structures with a specific distance separation was determined by the period T of the gas ejection from the disk surface. For example, we obtained a jet with a distance separation of about $11r_i$ associated with a period of 20 time units for the perturbation. However, for different values of T , an interesting effect occurs: the structures tend to dissipate at large axial distances for smaller values of T and to fragment into smaller structures for greater values of T .

Using this same numerical approach, Ouyed & Pudritz(1999) obtained episodic jets, using steady gas injection from the disk with velocities which were smaller than $10^{-3}v_{K,i}$ which we used. For injection velocities equal to this value, they obtained a highly collimated steady state jet⁵ with features similar to ours, with no perturbation in the velocity field of the disk. When we introduced perturbations with specific features in our steady state jet, our results showed the tendency to produce regular structures with separation distances $\sim 11r_i$, similar to the case where OP99 used a smaller velocity of $10^{-4}v_{K,i}$.

Gardiner & Frank (2000), using a magnetocentrifugal approach for the jet launching, studied an analytical one-dimensional model, in which they investigated the behaviour of a jet with a variable output speed. They inferred that a pulsing jet with an embedded helical magnetic field will inevitably develop a periodic field structure, consisting of rarefied regions (where the helix is “combed out”, resulting in a more poloidal geometry) alternating with denser, toroidally dominated knots. To confirm these predictions, they performed axisymmetric (2.5-dimensional) numerical simulations of a radiative MHD jet. They used conditions appropriate for protostellar jets and introduced a pulsed velocity with a sinusoidal

⁵They obtained steady state jets with velocities equal to or greater than $5 \times 10^{-4}v_{K,i}$, as well as episodic jets with different distances of separation for smaller velocities, using the same free parameters that we used.

form at the jet nozzle. Their results showed that pulsing jets with initially helical fields will evolve into an alternating poloidal-toroidal-dominated field geometry. The strongest toroidal field are confined to the dense knots, while predominantly poloidal fields are in the rarefied regions.

Both our results and those of Gardiner & Frank (2000) show that episodic jets are produced by introducing a pulsed mass load at the base of the jet. In particular, in our simulations, as well as in the study made by Ouyed & Pudritz (1997b), knot formation is associated with an anticorrelation of density and toroidal field strength. However, it should be noted that our study is concerned with the near-field region (i.e., the field region near the origin of the jet), whereas other studies, such as Gardiner & Frank (2000), are concerned with the far-field region.

There is currently some debate over the origin of the knots in protostellar flows. It remains unclear as to whether the knots are due to the pulsing of the source of the jet or to instabilities in the beam. Our results show that for a given constant ejection velocity from the disk surface which produces a steady state jet, an episodic jet is produced when the given ejection velocity is pulsed. We showed that the temporal and spatial knotty structure of the jets depends on the period of the velocity perturbation at the base of the jet.

Ouyed & Pudritz (1997b, 1999) rule out the possibility of a pinch MHD instability causing the knotty structure. This instability depends upon the strength of β and no such dependence was found by the authors. They found that only for β_i around unity, were knots produced. Birkinshaw (1990) noted that the Kelvin-Helmholtz (KH) instability does not exist if the beam speed is less than the Alfvén speed. Hardee & Rosen (1999) showed that KH growth rates increase as the magnetosonic Mach number decreases, provided the jet is super-Alfvénic and the magnetized jet is nearly stabilized to the KH instability when the jet is sub-Alfvénic. Hardee & Rosen (2002) found that the jets can be stabilized to the KH

helical and higher order asymmetric normal modes provided the velocity shear, $\Delta_u \equiv u_j - u_e$ between the jet and the external medium is less than a “surface” Alfvén speed, $v_{AS} \equiv [(\rho_J + \rho_E)(B_J^2 + B_E^2) / (4\pi\rho_J\rho_E)]^{1/2}$, where $\rho_J(\rho_E)$ is the jet (external medium) density and $B_J(B_E)$ is the jet (external medium) magnetic field. Ouyed et al. (2003) showed that the jet beyond the Alfvén surface becomes unstable to nonaxisymmetric KH instabilities, in three-dimensional simulations of jets from Keplerian accretion disks.

We also made a number of simulations using values of β_i different from unity. Figure 11 shows the results of one of these simulations, a periodic perturbation at the disk surface, using a period of $T = 20$. We plotted the isocontours of the density. The results for $\beta_i = 0.1$ are shown at the top of the figure. For this case, we see a smoothing of the structures along the jet. The isocontours of the density oscillate slightly near the jet axis. At the bottom of this figure, the results for $\beta_i = 10$ are shown. Here we observe the formation of structures from the jet axis up to a distance of about $z = 35r_i$. The region near the jet axis shows irregular structures and is less well defined (compare with Figure 5, where $\beta_i = 1$).

Our results are consistent with those of Hardee & Rosen (1999, 2002) and Ouyed et al. (2003). This is shown in the sequence of simulations in Figures 5 and 11. When β_i is small (strong magnetic fields, and the flow is sub-Alfvénic), there is a not much growth of the instability. The super-Alfvénic case in which β_i is close to unity (the flow speed is close to the Alfvén speed) is more unstable.

EDITOR: PLACE FIGURE 11 HERE.

We simulated a region very near the central object ($20r_i \times 80r_i$). It is interesting to evaluate the spatial dimensions involved in our simulations. For a protostar of mass $0.5M_\odot$, taking $r_i = 0.04$ AU for the inner radius of the disk, we have $80r_i \sim 10^{-5}$ pc. For a black hole of $10^8 M_\odot$ (the Schwarzschild radius is ~ 1 AU), assuming $r_i = 20.6$ AU, we have $80r_i \sim 10^{-2}$

pc. A distance of separation of $\sim 11r_i$ of the structures corresponds, for the protostar case, to $\sim 10^{-6}$ pc and for the black hole case, to $\sim 10^{-3}$ pc. These distances are much smaller than those between the Herbig-Haro objects observed in protostellar jets (about 10^{-2} pc). They are also much smaller than the distances observed between the emission knots, frequently found in AGN jets. In contrast to other simulations (e.g., Frank et al.(2000)), we obtain structures very near the formation region of the jets, involving distances not yet resolved by current instruments.

We can not say, at the present moment, whether or not the origin of the knotty structures seen in many jets at large distances from the source are due to the structures observed in the present simulations for regions close to the source. Further studies are needed to investigate this question.

The instabilities seen by Ouyed & Pudritz (1997b, 1999), producing the knotty structures, occur at the fast magnetosonic surface and propagate downstream. These instabilities, as explained by the authors, are due to MHD shocks. Some conditions in the knot generating region (in particular, for high $\mathbf{B}_\phi/\mathbf{B}_z$) seem to point towards a scenario in which the knot spacings could possibly be identified with the resonant wavelength pinch mode. As noted above, Ouyed & Pudritz (1997b, 1999) rule out the possibility of a pinch MHD instability causing the knotty structures since it has a strong β dependence, which was not found by the authors. The authors only observed knotty structures when β was around unity.

We believe that the knotty structures that we obtained in our simulations are due to the association of the pulsing mass load at the base of the jet and the instability due to MHD shocks, discussed in the studies of Ouyed & Pudritz (1997b, 1999). Our results indicate that a particular frequency of the pulsed velocity can excite the instability and produce persistent knotty structures along the jet.

The authors would like to thanks the editor, Steven Shore, and the anonymous referee for helpful comments. B. F. V. acknowledges the financial support of the Brazilian agency FAPESP under grant 95/99122. V. J. P. and R. O. would like to thank the Brazilian agency CNPq for partial support. We would like to thank the LCCA-USP for the possibility of using the CRAY J-90 computer in our simulations and especially the staff for their invaluable help. R. Ouyed and especially C. Fendt contributed to this paper with encouraging and valuable discussions, for which we are grateful. The authors also would like to thank the project PRONEX (No. 41.96.0908.00) for partial support. R. O. would like to thank the Brazilian agency FAPESP for partial support under grant 00/06770-2. V. J. P. would like to thank FAPESP for partial support under grant 2000/06769-4.

REFERENCES

- Bell, A.R., & Lucek, S.G. 1995, MNRAS, 277, 1327
- Birkinshaw, M. 1990, in Beams and Jets in Astrophysics, ed. P. A. Hughes (Cambridge: Cambridge Univ. Press)
- Chan, K.L., Henriksen R.N. 1980 ApJ, 241, 534
- Frank, A., Lery, T., Gardiner, T.A., Jones, T.W., & Ryu, D. 2000, ApJ, 540, 342 Goodson, A.P., Winglee, R.M., Böhm, K-H. 1997, ApJ, 489, 199
- Gardiner, T.A., Frank, A. 2000, ApJ, 545: L153-154
- Goodson, A.P., Winglee, R.M., & Böhm, K-H. 1997, ApJ, 489, 199
- Hardee, P.E, & Rosen, A. 1999, ApJ, 524, 650
- Hardee, P.E, & Rosen, A. 2002, astro-ph/0205377
- Hayashi, M., Shibatta, K., & Matsumoto, R. 1996, ApJ, 468:L37-40
- Krasnopolsky, R., Li, Z-Y., & Blandford, R. 1999, ApJ, 526, 631
- Königl, A., & Pudritz, R. 1999, Protostars and Planets IV, ed. by V. Manning, A. Boss, & S. Russel (U. Arizona Press:Tucson)
- Ouyed, R., & Pudritz, R.E. 1997a, ApJ, 482, 712 (OP97a)
- Ouyed, R., & Pudritz, R.E. 1997b ApJ, 484, 794
- Ouyed, R., & Pudritz, R.E. 1999 MNRAS, 309, 233
- Ouyed, R., Clarke, D.A., & Pudritz, R.E. 2003 ApJ, 582, 292
- Shibata, K., & Uchida, Y. 1986 PASJ, 38, 631

- Shu, F., Najita, H., Shang, Z., & Li, Z.-Y. 1999 *Protostars and Planets IV*, ed. by V. Manning, A. Boss, & S. Russel (U. Arizona Press: Tucson)
- Stone, J.M., & Norman, M.L. 1992a *ApJS*, 80, 753
- Stone, J.M., & Norman, M.L. 1992b *ApJS*, 80, 791
- Stone, J.M., & Norman, M.L. 1994 *ApJS*, 433, 746
- Ustyugova, G.V., Koldoba, A.V., Romanova, M.M., Chechetkin, V.M., & Lovelace, R.V.E. 1995 *ApJ*, 439, L39

Fig. 1.— Setup of the numerical simulations. Ghost zones ($z < 0$) define the surface of the Keplerian disk, while active zones (which are evolved in time) define the corona. The poloidal magnetic field continues smoothly into the disk surface (Fig. 2 of OP97a).

Fig. 2.— Initial conditions in the atmosphere: density distribution in units of ρ_i (eq. [8]) at the top and the magnetic potential (eq. [10]) at the bottom. Forty logarithmically spaced isocontour lines are shown for the density; 40 linearly spaced contours for the magnetic potential. The absolute nominal value for the steps between the isocontours of the logarithm of the density is 0.098. The velocity and toroidal magnetic field are initially zero.

Fig. 3.— Simulation A (without perturbation): (from top to bottom) Density, poloidal velocity, poloidal magnetic field, toroidal magnetic field and toroidal velocity for $\tau = 400$. Forty logarithmically spaced isocontour lines are shown for the density, and 40 linearly spaced contours for all the other quantities. The labels of the toroidal magnetic field and toroidal velocity denote 100 times their nominal values, in units of B_i and v_{Ki} , respectively. The absolute nominal values for the steps between the isocontours of the logarithm of the density, toroidal magnetic field and toroidal velocity are 0.1, $0.034B_i$ and $0.0118v_{Ki}$, respectively. The maximum value of the poloidal velocity is equal to $0.64v_{Ki}$. We note the formation of a steady state jet.

Fig. 4.— Simulation B ($T = 10$): (from top to bottom) Density, poloidal velocity, poloidal magnetic field, toroidal magnetic field and toroidal velocity for $\tau = 400$. Forty logarithmically spaced isocontour lines are shown for the density, and 40 linearly spaced contours for all the other quantities. The labels of the toroidal magnetic field and toroidal velocity denote 100 times their nominal values. The absolute nominal values for the steps between the isocontours of the logarithm of the density, toroidal magnetic field and toroidal velocity are 0.08, $0.0368B_i$ and $0.0108v_{Ki}$, respectively. The maximum value of the poloidal velocity is equal to $0.64v_{Ki}$. We note the formation of regularly spaced structures along the jet axis.

Fig. 5.— Simulation C ($T = 20$): (from top to bottom) Density, poloidal velocity, poloidal magnetic field, toroidal magnetic field and toroidal velocity for $\tau = 400$, using a better resolution (250, 80) uniform zones. Forty logarithmically spaced isocontour lines are shown for the density, and 40 linearly spaced contours for all the other quantities. The labels of the toroidal magnetic field and toroidal velocity denote 100 times their nominal values. The absolute nominal values for the steps between the isocontours of the logarithm of the density, toroidal magnetic field and toroidal velocity are 0.094, $0.0346B_i$ and $0.0154v_{Ki}$, respectively. The maximum value of the poloidal velocity is equal to $0.61v_{Ki}$. We note, again, the formation of regularly spaced structures along the jet axis.

Fig. 6.— Simulation D ($T = 40$): (from top to bottom) Density, poloidal velocity, poloidal magnetic field, toroidal magnetic field and toroidal velocity for $\tau = 400$. Forty logarithmically spaced isocontour lines are shown for the density, and 40 linearly spaced contours for all the other quantities. The labels of the toroidal magnetic field and toroidal velocity denote 100 times their nominal values. The absolute nominal values for the steps between the isocontours of the logarithm of the density, toroidal magnetic field and toroidal velocity are 0.094, $0.0342B_i$ and $0.016v_{Ki}$, respectively. The maximum value of the poloidal velocity is equal to $0.67v_{Ki}$. We note fragmentation of regularly spaced structures along the jet axis.

Fig. 7.— Simulation E ($T = 80$): (from top to bottom) Density, poloidal velocity, poloidal magnetic field, toroidal magnetic field and toroidal velocity for $\tau = 400$. Forty logarithmically spaced isocontour lines are shown for the density, and 40 linearly spaced contours for all the other quantities. The labels of the toroidal magnetic field and toroidal velocity denote 100 times their nominal values. The absolute nominal values for the steps between the isocontours of the logarithm of the density, toroidal magnetic field and toroidal velocity are 0.096, $0.0344B_i$ and $0.0152v_{Ki}$, respectively. The maximum value of the poloidal velocity is equal to $0.64v_{Ki}$. We note more pronounced fragmentation of the regularly spaced structures

along the jet axis.

Fig. 8.— Simulations A to E: Density (solid line), axial velocity (dotted line), toroidal velocity (dashed line) and toroidal magnetic field (dashed-dotted line) along the z -axis at $r = 5r_i$. The curves are normalized by their maximum values.

Fig. 9.— Simulations A to E: Axial and radial mass fluxes (eq.[18]). Here we show the axial (solid line) and radial (dotted line) mass fluxes at $\tau = 400$. We calculate the axial mass flux, at each axial distance z , using the transverse area $A = \pi r^2$ with $r = 20r_i$. For the radial mass flux, we use the cylindrical surface area $A = 2\pi r z$ with $r = 20r_i$, at each axial distance z . The fluxes are given in units of $\dot{m}_i = 2\pi r_i^2 \rho_i v_{Ki}$. The scale on the vertical axis represents a tenth of the nominal value.

Fig. 10.— Simulations A to E: The sonic Mach numbers (dotted lines), Alfvénic Mach numbers (dashed-dotted lines) and magnetosonic Mach numbers (solid lines) along the z -axis for all simulations at $\tau = 400$ in a cut at $r = 5r_i$. For simulations C to E, dashed-dotted lines denote 10^{-2} times the nominal values of Alfvénic Mach numbers.

Fig. 11.— Logarithmically spaced isocontour lines are shown for the density. On the top we have the result $\beta_i = 0.1$. We obtained, in this case, smoothing of the structures along the jet. The isocontours of the density slightly oscillate near the jet axis. On the bottom of the figure we used $\beta_i = 10$. Here we observe the formation of structures displaced from the jet axis up to the distance of about $z = 35r_i$. The region near the jet axis show irregular structures and is less well defined.